



British Mycological
Society promoting fungal science

journal homepage: www.elsevier.com/locate/funbio



Species of Mycosphaerellaceae and Teratosphaeriaceae on native Myrtaceae in Uruguay: evidence of fungal host jumps

C. A. PÉREZ^{a,*}, M. J. WINGFIELD^b, N. ALTIER^c, R. A. BLANCHETTE^d

^aDepartamento de Protección Vegetal, Facultad de Agronomía, Universidad de la República, Uruguay

^bForestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, South Africa

^cInstituto Nacional de Investigación Agropecuaria (INIA), Uruguay

^dDepartment of Plant Pathology, University of Minnesota, USA

ARTICLE INFO

Article history:

Received 25 July 2012

Received in revised form

1 December 2012

Accepted 6 December 2012

Available online 21 December 2012

Corresponding Editor:

H. Thorsten Lumbsch

Keywords:

Host shift

Mycosphaerella Leaf Disease

Myrtaceae diseases

ABSTRACT

Mycosphaerella species are well-known causal agents of leaf diseases on many economically and ecologically important plant species. In Uruguay, a relatively large number of *Mycosphaerellaceae* and *Teratosphaeriaceae* are found on *Eucalyptus*, but nothing is known of these fungi on native *Myrtaceae*. The aim of this study was to identify *Mycosphaerellaceae* and *Teratosphaeriaceae* species associated with leaf diseases on native *Myrtaceae* in Uruguay and to consider whether host jumps by the pathogen from introduced *Eucalyptus* to native *Myrtaceae* have occurred. Several native forests throughout the country were surveyed with special attention given to those located close to *Eucalyptus* plantations. Five species belonging to the *Mycosphaerellaceae* and *Teratosphaeriaceae* clades were found on native *Myrtaceous* trees and three of these had previously been reported on *Eucalyptus* in Uruguay. Those occurring both on *Eucalyptus* and native *Myrtaceae* included *Pallidocercospora heimii*, *Pseudocercospora norchiensis*, and *Teratosphaeria aurantia*. In addition, *Mycosphaerella yunnanensis*, a species known to occur on *Eucalyptus* but not previously recorded in Uruguay, was found on leaves of two native *Myrtaceous* hosts. Because most of these species occur on *Eucalyptus* in countries other than Uruguay, it appears that they were introduced in this country and have adapted to be able to infect native *Myrtaceae*. These apparent host jumps have the potential to result in serious disease problems and they should be carefully monitored.

© 2012 The British Mycological Society. Published by Elsevier Ltd. All rights reserved.

Introduction

A diverse group of *Mycosphaerellaceae* and *Teratosphaeriaceae* has been associated with *Mycosphaerella* leaf disease (MLD), which are considered particularly important in *Eucalyptus* plantations worldwide (Park *et al.* 2000; Maxwell *et al.* 2003; Crous *et al.* 2004; Crous *et al.* 2006; Cortinas *et al.* 2006; Hunter *et al.* 2006; Summerell *et al.* 2006; Crous *et al.* 2009; Perez *et al.*

2009a; Hunter *et al.* 2011). These fungi cause leaf spots, leaf blotches, or petiole and stem cankers that often result in stressed and stunted trees, adversely affecting commercial forestry operations (Lundquist & Purnell 1987; Carnegie *et al.* 1994; Carnegie *et al.* 1998; Park *et al.* 2000; Sánchez Márquez *et al.* 2011).

Although most studies on MLD have focused primarily on *Eucalyptus*, species of *Mycosphaerellaceae* and *Teratosphaeriaceae*

* Corresponding author. Estacion Experimental "Dr. Mario Cassinoni", Facultad de Agronomía, Universidad de la República, Ruta 3, km 363, Paysandu, CP 60000, Uruguay. Tel.: +598 47227950; fax: +598 47227950.

E-mail address: caperez@fagro.edu.uy (C. A. Pérez).

1878-6146/\$ – see front matter © 2012 The British Mycological Society. Published by Elsevier Ltd. All rights reserved.

<http://dx.doi.org/10.1016/j.funbio.2012.12.002>

have also been found infecting species of the Myrtaceae other than those on *Eucalyptus* genus. At least 23 species have been found on non-*Eucalyptus* species in the Myrtaceae worldwide (Crous 1999; Sivanesan & Shivas 2002; Carnegie *et al.* 2007). Increased sampling over the last decade has shown that certain species included in these taxa can be found on a wide range of different hosts including different plant orders such as Myrtales, Proteales, Fabales and Apiales (Hunter *et al.* 2011). The apparent ease of movement in these fungi between hosts is thus of considerable concern.

Most *Eucalyptus* species are native to Australia and have been moved extensively around the world. Where *Eucalyptus* spp. are grown as non-natives, they have largely been separated from their natural enemies (Burgess & Wingfield 2002; Wingfield 2003). This is a situation that is gradually changing with pathogens and pests being brought back into contact with their hosts due to accidental introductions resulting in serious disease problems (Wingfield *et al.* 2008).

Eucalypts are threatened not only by pathogens that are known to attack them in their native environment but there is also growing evidence of pathogens from native Myrtaceae undergoing host shifts to infect them (Slippers *et al.* 2005). The best-known example of such a host shift linked to *Eucalyptus* is that of the *Eucalyptus* rust pathogen *Puccinia psidii*. This rust disease is native on Myrtaceae in South and Central America and has adapted to infect *Eucalyptus* in that region (Coutinho *et al.* 1998; Glen *et al.* 2007). In addition, there are many recent examples of members of the Cryphonectriaceae, that are native on members of the Myrtales, adapted to infect *Eucalyptus* in Africa (Heath *et al.* 2006) as well as South and Central America and Asia (Hodges *et al.* 1986; Myburg *et al.* 2003; Rodas *et al.* 2005; Gryzenhout *et al.* 2006).

Where pathogens have been introduced into new areas, they also have the potential to cause serious diseases on related native plants. It is for this reason that the severe impact of the recent introduction of *P. psidii* to Australia has been of such great concern (Grgurinovic *et al.* 2006; Glen *et al.* 2007; Carnegie *et al.* 2010).

Eucalyptus is widely planted in Uruguay and these trees have already been seriously affected by many diseases thought to have been introduced from other areas. Yet almost nothing is known regarding the pathogens of native Myrtaceae in Uruguay, whether these trees might be threatened by *Eucalyptus* pathogens or possibly the pathogens on the Myrtaceae causing new disease problems on *Eucalyptus*. Uruguay has a large resource of native Myrtaceae (Brussa & Grela 2007) and the aim of this study was to identify Mycosphaerellaceae and Teratosphaeriaceae species associated with MLD on native Myrtaceae species. Furthermore, we considered their relationships with those species currently affecting *Eucalyptus* plantations in Uruguay.

Materials and methods

Samples and isolations

Between 2005 and 2008, trees belonging to the Myrtaceae were surveyed in native forests throughout Uruguay and special attention was placed on those located close to *Eucalyptus*

plantations. Leaves showing MLD symptoms were recorded photographically, collected and taken to the laboratory for further study. Samples were collected from a total of 199 trees belonging to 20 native species residing in the Myrtaceae (Table 1). Sampled trees were distributed over the main areas where *Eucalyptus* is planted (Fig 1).

Lesions on leaves bearing pseudothecia were processed for isolation following the procedure described by Crous (1998). Parts of lesions with mature pseudothecia were soaked in sterile water for 2 h. The leaf pieces were then dried on sterilized paper and adhered with adhesive tape to the undersides of Petri dish lids with the pseudothecia facing the surface of 2 % malt extract agar (MEA) (2 % malt extract, 1.5 % agar; Oxoid, Basingstoke, England). Petri dishes were incubated at 17–18 °C in the dark for 24–48 h. Ascospores that had been ejected onto the media and had germinated were observed under a microscope to record the germination patterns as described by Crous (1998). Individual germinating ascospores were lifted from the medium and transferred to new plates to generate monosporic cultures.

Where pseudothecia were not observed, pieces of leaf from the edges of the lesion were cut, surface-disinfested in 70 % ethyl alcohol for 30 s, and rinsed twice in sterile distilled water, blotted dry on sterile filter paper, and plated on 2 % MEA amended with 0.01 g of streptomycin per litre to minimize bacterial contamination. Plates were then incubated at room temperature and emerging colonies were sub-cultured on fresh 2 % MEA plates. Only those cultures with colony morphologies resembling those of species of Mycosphaerellaceae/Teratosphaeriaceae were included in further studies.

Table 1 – List of Myrtaceae species native to Uruguay sampled in this study. Tree species in bold indicate those where Mycosphaerellaceae or Teratosphaeriaceae species were found associated with MLD symptoms.

Tree Species	Fungal species	Province
<i>Acca sellowiana</i> *	<i>Pseudocercospora norchiensis</i> <i>Passalora loranthi</i>	Rivera Rivera
<i>Agariota eucalyptides</i>		
<i>Blepharocalyx salicifolius</i>*	<i>Mycosphaerella yunnanensis</i> <i>Pseudocercospora norchiensis</i> <i>Teratosphaeria aurantia</i>	Rivera Rivera Rivera
<i>Calyptanthes concinna</i>		
<i>Eugenia involucrata</i>		
<i>E. mansonii</i>		
<i>E. repanda</i>		
<i>E. uniflora</i>		
<i>E. uruguayensis</i>		
<i>Gomidesia palustris</i>		
<i>Hexachlamis edulis</i>		
<i>Myrceugenia euosma</i>		
<i>Myrce. glaucescens</i>*	<i>Pallidocercospora heimii</i>	Río Negro
<i>Myrcianthes cisplatensis</i>		
<i>Myrci. pungens</i>		
<i>Myrciaria tenella</i>		
<i>Myrrhinium atropurpureum</i> var. <i>octandrum</i>*	<i>Mycosphaerella yunnanensis</i>	Rivera
<i>Psidium luridum</i>		
<i>P. incanum</i>		
<i>P. pubifolium</i>		




Fig 1 – Map of Uruguay. The shaded area indicates the geographic distribution of *Eucalyptus* plantations in the country. Stars indicate those locations where *Mycosphaerellaceae* and *Teratosphaeriaceae* species were found occurring on native trees.

For these isolates, pure cultures were made by transferring hyphal tips to clean culture media and thus ensuring that isolates represented a single genotype. Cultures were grouped based on ascospore germination pattern, conidial and ascospore morphology, and colony morphology. These morphological characteristics were then used to confirm grouping that emerged from the phylogenetic analyses.

DNA extraction, polymerase chain reaction (PCR), sequencing, and phylogenetic analysis

DNA was extracted from isolates representing each morphological group. Mycelium was scrapped directly from the surface of colonies grown on 2 % MEA plates at room temperature for 30 d and transferred to Eppendorf tubes (1.5 ml) with 3-mm glass beads and extraction buffer (Qiagen Inc., Valencia, CA, USA). These were shaken vigorously using a vortex mixer and placed in a water bath at 60 °C for 1 h. DNA extraction was performed using the Qiagen Plant DNeasy Mini Kit (Qiagen Inc., Valencia, CA, USA) following manufacturer's instructions.

The entire ribosomal DNA internal transcribed spacer (ITS) regions (ITS1 and ITS2) including the 5.8S gene of the rDNA were amplified using the primers ITS1 and ITS4 (White et al. 1990). PCR amplifications were performed in an MJ Research PTC 200 DNA Engine Thermal Cycler PCR (MJ Research, Reno, NV, USA) with the following conditions: initial

denaturation for 5 min at 94 °C, then 35 cycles of 1 min at 94 °C, 1 min at 50 °C, 1 min at 72 °C, followed by a final elongation step of 5 min at 72 °C, and hold at 10 °C. The PCRs had a total volume of 25- μ l containing 1X of Amplitaq Gold PCR Master-Mix (Applied Biosystems, Foster City, CA, USA), 0.2 μ M of each primer and approx. 10 ng μ l⁻¹ of DNA template. Deionized-distilled water was added to a final volume of 25 μ l.

PCR products were stained with SYBR Green nucleic acid dye (MBL International, Woburn, MA, USA) and visualized on 1.5 % agarose gels under UV light. Amplicons were prepared for sequencing using ExoSAP-IT PCR clean-up kit (USB Corp., Cleveland, OH, USA) following the manufacturer's instructions. For sequencing reactions, the same primer pairs were used with the ABI Prism Dye Terminator Cycle Sequencing Ready Reaction Kit (Applied Biosystems, Foster City, CA, USA) and an ABI Prism 377 automated DNA sequencer. Forward and reverse sequences were assembled using ChromasPro software version 1.33 (Technelysium Pty. Ltd., Eden Prairie, MN, USA).

Sequences were subjected to BLAST searches in NCBI Genbank (<http://www.ncbi.nlm.nih.gov/blast/Blast.cgi>, verified 26 Jun. 2012), and those of the most closely matching species were downloaded from GenBank. Where available, sequences that represented the ex-type cultures of the closely matching species were used and all species of *Mycosphaerellaceae* and *Teratosphaeriaceae* previously reported from *Eucalyptus* were also included. Following a first preliminary phylogenetic analysis, the alignment was trimmed, discarding those species only distantly related to the sequences under investigation and populating the remainder of the data set with at least two sequences per taxon when possible (Table 2). In addition, sequences of species residing in the *Mycosphaerellaceae* and *Teratosphaeriaceae* obtained from *Eucalyptus* spp. in Uruguay (Perez et al. 2009b) were included in the alignment for comparison. Multiple sequence alignments were made online using the E-INS-i strategy in MAFFT version 6 (<http://align.bmr.kyushu-u.ac.jp/mafft/online/server/>, verified 4 May 2012) (Kato et al. 2005).

Neighbour-joining (NJ) and Maximum parsimony (MP) analyses were performed using PAUP v. 4.0b10 (Swofford 2002). The best substitution model for NJ analysis was determined using Modeltest v. 3.7 (Posada & Crandall 1998) from which a general time reversible substitution model including a proportion of invariant sites and gamma-distributed substitution rates of the remaining sites (GTR + I + G) was selected from the Akaike information criterion (AIC; proportion of invariable sites (I) = 0.3375; gamma distribution shape parameter (G) = 0.8871; base frequencies: $\pi_A = 0.2015$, $\pi_C = 0.2991$, $\pi_G = 0.2800$, $\pi_T = 0.2194$). Gaps were treated as missing data and all characters were treated as unordered and of equal weight. MP analysis was performed using the heuristic search option with simple addition of taxa and tree bisection and reconnection (TBR) as the branch-swapping algorithm. Support for the nodes of the shortest trees was determined by analyses of 1000 bootstrap replicates (Hillis & Bull 1993) and tree length (TL), consistency index (CI), retention index (RI), and homoplasy index (HI) were calculated.

Part of the translation elongation factor 1- α (EF1- α) region and the Actin (ACT) gene was also amplified for selected isolates to conduct a multiple gene genealogy analysis and thus

Table 2 – List of sequences used in the phylogenetic analysis including those obtained in this study and reference sequences obtained from Genbank. Cultures from Uruguay are indicated with the prefix ‘UY’. Cultures from native Myrtaceae sequenced in this study are in bold.

Culture ID	Species	Host	GenBank accession no.		
			ITS	EF-1 α	ACT
UY322	<i>Pallidocercospora heimii</i>	<i>Myrceugenia glaucescens</i>	EU853466	JX444995	JX445002
UY372	<i>Teratosphaeria aurantia</i>	<i>Eucalyptus grandis</i>	EU851913		
UY423	<i>Pal. heimii</i>	<i>Euc. dunnii</i>	EU851921	JX444996	JX445003
UY1382	<i>T. aurantia</i>	<i>B. salicifolius</i>	EU853472	JX444997	JX445004
UY1436	<i>Pseudocercospora norchiensis</i>	<i>Acca sellowiana</i>	EU853474	JX444998	JX445005
UY1462	<i>Mycosphaerella yunnanensis</i>	<i>Myrrhimum atropurpureum</i> var. <i>octandrum</i>	EU853475		
UY1483	<i>M. yunnanensis</i>	<i>B. salicifolius</i>	EU853477		
UY1484	<i>Ps. norchiensis</i>	<i>B. salicifolius</i>	EU853478	JX444999	JX445006
UY1506	<i>Passalora loranthi</i>	<i>Acca sellowiana</i>	EU853479		
UY1528	<i>Ps. norchiensis</i>	<i>Euc. dunnii</i>	EU851936	JX445000	JX445007
UY1561	<i>Ps. norchiensis</i>	<i>Euc. grandis</i>	EU851938	JX445001	JX445008
CBS120735 T	<i>M. elongata</i>	<i>E. camaldulensis</i> × <i>E. urophylla</i>	EF394833		
CBS111519 T	<i>M. endophytica</i>	<i>Eucalyptus</i> sp.	DQ267579		
CBS111001 T	<i>M. keniensis</i>	<i>Euc. grandis</i>	AF309601		
CBS326.52	<i>'M'. laricina</i>	<i>Larix decidua</i>	AY152590		
Lari01.03	<i>'M'. laricina</i>	n/a	DQ019342		
CBS118493 T	<i>M. scytalidii</i>	<i>Eucalyptus</i> sp.	DQ303016		
STE-U2769	<i>M. walkeri</i>	<i>Eucalyptus</i> sp.	AF309616		
CBS119975 T	<i>M. yunnanensis</i>	<i>Eucalyptus</i> sp.	DQ632686		
CBS119976	<i>M. yunnanensis</i>	<i>Eucalyptus</i> sp.	DQ632687		
CMW23445	<i>M. yunnanensis</i>	<i>Eucalyptus</i> sp.	DQ632688		
CPC3837 T	<i>Pal. acaciigena</i>	<i>Acacia mangium</i>	AY752143		
CBS681.95 T	<i>Pal. crystallina</i>	<i>Euc. bicostata</i>	AY490757		
CBS110682 T	<i>Pal. heimii</i>	<i>Eucalyptus</i> sp.	DQ239992	DQ211667	DQ147638
CBS120743	<i>Pal. heimii</i>	<i>Euc. urophylla</i>	EF394838		
CMW5719	<i>Pal. heimii</i>	<i>Eucalyptus</i> sp.	AF452516		
CPC13371	<i>Pal. heimii</i>	<i>Euc. urophylla</i>	EF394840		
CBS111190 T	<i>Pal. heimioides</i>	<i>Eucalyptus</i> sp.	AF309609		
CBS111364	<i>Pal. heimioides</i>	<i>Eucalyptus</i> sp.	DQ267586		
CBS114774 T	<i>Pal. irregulariramosa</i>	<i>Euc. saligna</i>	AF309607		
STE-U2123	<i>Pal. konae</i>	<i>Leucadendron</i> sp.	AY260086		
STE-U2125	<i>Pal. konae</i>	<i>Leucadendron</i> sp.	AY260085		
n/a	<i>Pas. loranthi</i>	n/a	AY348311		
CPC11258	<i>Pas. sequoiae</i>	<i>Juniperus virginiana</i>	GU214667		
CMW5148 T	<i>Ps. basiramifera</i>	<i>Euc. pellita</i>	AF309595		
CBS111280	<i>Ps. basitruncata</i>	<i>Euc. grandis</i>	DQ267601	DQ211676	DQ147621
CBS114664	<i>Ps. basitruncata</i>	<i>Euc. grandis</i>	DQ267600		
CBS110969 T	<i>'Ps.' colombiensis</i>	<i>Euc. urophylla</i>	AY752149		
CBS682.95 T	<i>'Ps.' epispermogonia</i>	<i>Euc. grandis</i>	DQ267587		
CBS110777 T	<i>Ps. eucalyptorum</i>	<i>Eucalyptus</i> sp.	AF309598		
CMW13586 T	<i>Ps. flavomarginata</i>	<i>Euc. camaldulensis</i>	DQ155657		
STE-U2556	<i>Ps. luzardii</i>	<i>Hancornia speciosa</i>	AF362057		
CBS111069 T	<i>Ps. natalensis</i>	<i>Eucalyptus</i> sp.	DQ303077		
CBS120738 T	<i>Ps. norchiensis</i>	<i>Eucalyptus</i> sp.	EF394859		
STE-U1458	<i>Ps. paraguayensis</i>	<i>Eucalyptus</i> sp.	AF309596		
CBS120029 T	<i>Ps. schizolobii</i>	<i>Schizolobium parahybum</i>	DQ885903		
CPC10547 T	<i>'Ps.' thailandica</i>	<i>Acacia mangium</i>	AY752156	AY840478	AY752219
CBS680.95 T	<i>Teratosphaeria africana</i>	<i>Euc. viminalis</i>	AF309602	DQ235099	DQ147609
CMW3025	<i>T. africana</i>	<i>Euc. viminalis</i>	AF283690		
CBS110500 T	<i>T. aurantia</i>	<i>Euc. globulus</i>	AY725531	DQ235097	DQ147610
MURU151	<i>T. aurantia</i>	<i>Euc. globulus</i>	AY150331		
MURU152	<i>T. aurantia</i>	<i>Euc. globulus</i>	AY509742		
MURU222	<i>T. aurantia</i>	<i>Euc. globulus</i>	AY509744		
CBS120146 T	<i>T. molleriana</i>	<i>Eucalyptus</i> sp.	EF394844		
CBS116005 T	<i>T. nubilosa</i>	<i>Euc. globulus</i>	AF309618		
CBS110949 T	<i>T. ohnowa</i>	<i>Euc. grandis</i>	AY725575		
CBS118508 T	<i>T. pluritubularis</i>	<i>Euc. globulus</i>	DQ303007		
CBS120303 T	<i>T. gauchensis</i>	<i>Euc. grandis</i>	EU019290		
CBS114238 T	<i>Uwebraunia commune</i>	<i>Euc. globulus</i>	AY725541		
CBS110748 T	<i>U. dekkeri</i>	<i>Eucalyptus</i> sp.	AF309624		

(continued on next page)

Table 2 – (continued)

Culture ID	Species	Host	GenBank accession no.		
			ITS	EF-1 α	ACT
STE-U348	<i>Zasmidium marasasii</i>	<i>Syzygium</i> sp.	AF309591		
CMW3358 T	<i>Zasmidium parkii</i>	<i>Eucalyptus</i> sp.	AF309590		
CMW7773	<i>Neofusicoccum ribis</i>	<i>Ribes</i> sp.	AY236936		

^T: ex-type cultures.

to confirm the identity of those species found on both native Myrtaceae and *Eucalyptus*. EF-1 α was amplified using the primers EF1-728F (5'-CAT CGA GAA GTT CGA GAA GG-3') and EF1-986R (5'-TAC TTG AAG GAA CCC TTA CC-3') (Carbone & Kohn 1999). Reaction conditions were: an initial denaturation step of 96 °C for 2 min, followed by 35 cycles of denaturation at 94 °C for 30 s, primer annealing at 56 °C for 30 s and primer extension at 72 °C for 30 s. The reaction was completed with a final extension at 72 °C for 7 min. For the ACT gene, the primers ACT-512F (5'-ATG TGC AAG GCC GGT TTC GC-3') and ACT-783R (5'-TAC GAG TCC TTC TGG CCC AT-3') were used (Carbone & Kohn 1999). PCR reaction conditions were: an initial denaturation step at 96 °C for 2 min, followed by 10 cycles of denaturation at 94 °C for 30 s, primer annealing at 61 °C for 45 s and extension at 72 °C for 45 s. This was followed by 25 cycles of denaturation at 94 °C for 30 s, primer annealing at 61 °C and elongation at 72 °C for 45 s with an increase of 5 s per cycle. The reaction was completed with a final elongation step at 72 °C for 7 min.

ITS, EF1- α , and ACT sequence datasets were examined for congruence using the Partition Homogeneity Test in PAUP (Farris et al. 1995; Huelsenbeck et al. 1996). Selected isolates of *Pallidocercospora heimii*, *Pseudocercospora norchiensis* and *Teratosphaeria aurantia*, obtained from *Eucalyptus* (Perez et al. 2009b), were included as references to confirm genetic similarity. A combined data set for the ITS, EF1- α , and ACT sequence was analyzed using NJ and MP. Phylogenetic analyses were performed using PAUP Version 4.0b10. Best models for NJ analysis was determined from the AIC Modeltest version 3.7 (Posada & Crandall 1998) as GTR + G for the combined dataset (proportion of invariable sites (I) = 0; gamma distribution shape parameter (G) = 0.2668; base frequencies: π_A = 0.2136, π_C = 0.2966, π_G = 0.2583, π_T = 0.2315). Gaps generated in the alignment process during the comparison were treated as missing data and all characters were treated as unordered and of equal weight. Ties were broken randomly when found.

All the sequences obtained in this study were deposited in GenBank (Table 1). Sequence alignments and trees of the ITS alone and combined analysis have been deposited in TreeBASE (accession number: 13107).

Results

Samples and isolations

Twenty species of Myrtaceae from a wide range of different collection sites in Uruguay were evaluated and sampled during this study. Symptoms resembling MLD were observed on four species namely *Acca sellowiana*,

Blepharocalyx salicifolius, *Myrceugenia glaucescens* and *Myrrhinium atropurpureum* var. *octandrum* (Table 1). A total of 45 isolates were obtained from lesions on leaves of these trees. Isolates were grouped by culture and conidial morphology, ascospore germination pattern and host species. One isolate was then selected from each of the resulting seven groups for further investigation using DNA sequence comparisons.

DNA comparison and phylogenetic analyzes

ITS analysis

ITS sequences were generated in both directions and DNA amplicons of ~550 nucleotides were obtained after assemblage. Sequences were deposited in GenBank and accession numbers are shown in Table 2. Following BLAST searches, the 11 sequences obtained in this study were aligned with closest taxa available in GenBank or other species of Mycosphaerellaceae and Teratosphaeriaceae that have been previously reported on *Eucalyptus*. The alignment consisted of 63 ingroup sequences and *Neofusicoccum ribis* as the outgroup taxon.

Sequence alignment resulted in a total of 708 characters of which 233 were constant, 111 variable characters were parsimony-uninformative and 364 were parsimony informative. NJ and MP resulted in trees of identical topology. The heuristic search analysis of the data resulted in four most parsimonious trees (TL = 1129 steps; CI = 0.648; RI = 0.841; HI = 0.352). The genetic distance consensus tree obtained with NJ analysis based on the ITS region of the rDNA operon analysis is shown (Fig 2) with the bootstrap values of 1000 replicates of NJ and MP analyses displayed on the branches.

Combined data set analysis (ITS, EF-1 α , ACT)

The five sequences of those species previously found on *Eucalyptus* in Uruguay were subject to a combined analysis to fully verify their identity. The sequence alignment consisted of 12 ingroup sequences and *Teratosphaeria nubilosa* as the outgroup. A combined dataset of a total of 984 characters was analyzed (474, 227, 283 characters for ITS, EF-1 α , and ACT, respectively), of which 532 were constant, 147 were parsimony-uninformative and 305 were parsimony informative. Both NJ (NJ) and MP analyses resulted in trees of identical topology. Heuristic search analysis of the data resulted in one most parsimonious tree (TL = 811 steps; CI = 0.874; RI = 0.905; HI = 0.126). The distance tree obtained from NJ analysis of the combined ITS, EF1- α , and ACT with the bootstrap values of 1000 replicates from NJ and MP analyses displayed on the branches is shown (Fig 3).




Fig 2 – Genetic distance consensus tree obtained with NJ analysis based on the ITS region of the rDNA operon. Species name and GenBank accession number are shown for each sequence. Sequences labelled with a ‘T’ at the end correspond to the ex-type culture. Bootstrap values of 1000 replicates of NJ and MP analyses are shown above and below branches, respectively. Only bootstrap values higher than 75 % are shown. *Neofusicoccum ribis* was used as outgroup taxon. Uruguayan isolates are indicated with the prefix ‘UY’ and sequences corresponding to isolates obtained from native Myrtaceous trees are in bold. Underlined sequences correspond to isolates obtained from *Eucalyptus* spp. in Uruguay, and included for reference.




Fig 3 – NJ tree from the combined analysis based on partial ITS, Actin, and EF1- α sequence data. Bootstrap values of 1000 replicates of NJ and MP analyses are shown above and below branches, respectively. Only bootstrap values higher than 75 % are shown. Sequences corresponding to isolates obtained from native Myrtaceous trees are in bold.

Species identified

A diverse group of Mycosphaerellaceae and Teratosphaeriaceae were found to occur on diseased leaves of native Myrtaceae in this study. Phylogenetic analyses revealed a total of five distinct species residing in the Mycosphaerellaceae and Teratosphaeriaceae. These included *Mycosphaerella yunnanensis*, *Pallidocercospora heimii*, *Passalora lorantheri*, *Pseudocercospora norchiensis*, and *Teratosphaeria aurantia*.

Three of these species were previously found on *Eucalyptus* by Pérez et al. (2009b). *Teratosphaeria aurantia* was found associated with leaf spots on *Blepharocalyx salicifolius*, in the northern region (i.e. province of Rivera). The sequence of this isolate was identical to that of UY372 obtained from *Eucalyptus grandis* by Pérez et al. (2009a,b,c) and was confirmed in the multi-gene analysis.

Pallidocercospora heimii was associated with leaf spots only on *Myrceugenia glaucescens* in Rio Negro, and the sequence of this isolate was identical to that for the isolate obtained from *Eucalyptus dunnii* by Pérez et al. (2009b), whereas *Ps. norchiensis* was found on leaves of two native tree species, *Acca sellowiana* and *B. salicifolius*, also in the province of Rivera, and obtained sequences grouped with sequences UY1528 and UY1561 obtained from *E. dunnii* and *E. grandis* also in Rivera by Pérez et al. (2009b). *Passalora lorantheri* was found on leaf spots on *A. sellowiana* in the province of Rivera. Surprisingly, *M. yunnanensis*, not previously reported in Uruguay, was found on leaves of *B. salicifolius* and *Myrrhinium atropurpureum* var. *octandrum* in Rivera. Phylogenetic grouping was strongly supported by both NJ and MP analyses and sequences differed only at one base from the sequence for the ex-type culture of this fungus.

Discussion

Results of this study clearly showed that there is a relatively diverse group of species belonging to the Mycosphaerellaceae and Teratosphaeriaceae associated with leaf spots on native Myrtaceae in Uruguay. Three species, *Pallidocercospora heimii*, *Pseudocercospora norchiensis* and *Teratosphaeria aurantia*, are well-known *Eucalyptus* leaf spot associated fungi, previously reported to infect *Eucalyptus* in Uruguay (Balmelli et al. 2004; Crous et al. 2006; Pérez et al. 2009b). An important and intriguing aspect of this study was the clear evidence of fungi previously thought to be specific to *Eucalyptus*, occurring on the leaves of native trees in Uruguay. These fungi are all known on *Eucalyptus* leaves in countries other than Uruguay and it seems most likely that they were introduced into Uruguay on *Eucalyptus* and have subsequently undergone a host shift to native tree species. Such host shifts have recently been shown in Uruguay for *Quambalaria leaf disease* caused by *Quambalaria eucalypti* (Pérez et al. 2008) and *Neofusicoccum eucalyptorum* (Pérez et al. 2009c). This is, however, the first evidence of species associated with MLD of *Eucalyptus* undergoing such host shifts.

Teratosphaeria aurantia (syn. *Mycosphaerella aurantia*) was found associated with leaf spots on *Blepharocalyx salicifolius*. Although there has been some confusion regarding the identification of this species, with Hunter et al. (2006) suggesting that it is likely the same as *Teratosphaeria africana*, isolate UY1382 obtained from *B. salicifolius* grouped strongly with other *T. aurantia* isolates, including that of the ex-type (Maxwell et al. 2003). Morphological characteristics of ascospores observed in this study also showed they did not have constrictions at the median septum, as described for *T. aurantia* (Maxwell et al. 2003). To date, this species has only been known to occur in Australia and Uruguay (Andjic et al. 2010; Hunter et al. 2011; Pérez et al. 2009b) and this study represents the first report on a host other than *Eucalyptus*.

Pallidocercospora heimii was found associated with leaf lesions on *Myrceugenia glaucescens*. *Pallidocercospora* is a novel genus to accommodate former '*Mycosphaerella*' species with pale brown Cercospora-like conidia, and includes *Pal. heimii* (syn. *Mycosphaerella heimii*) (Crous et al. 2013). Although Hunter et al. (2006) considered *Pal. heimii* to represent a member of a species complex due to the difficulty differentiating this species from *Pallidocercospora heiminoidea*, *Pallidocercospora crystallina* and *Pallidocercospora irregulariramosa*, we found that isolate UY322 consistently grouped with *Pal. heimii* sequences and it was clearly separate from other related species of this complex. The ITS sequence for this isolate was also identical to isolate UY423 obtained from *Eucalyptus dunnii* in Uruguay by Pérez et al. (2009b).

Pallidocercospora heimii is known from Australia, Brazil, Madagascar, Portugal, Uruguay, and Venezuela (Hunter et al. 2004; Crous et al. 2006; Crous et al. 2007; Pérez et al. 2009b) where it has been found only on *Eucalyptus*. However, it was also found on *Acacia auriculiformis* and *Acacia* sp. in Thailand (Crous & Groenewald 2005). Our results support this data and demonstrate that it is able to cross hosts and all indications are that in Uruguay, it has moved from *Eucalyptus* onto native Myrtaceae. *Pseudocercospora norchiensis*,

found on *Acca sellowiana* and *B. salicifolius* in this study, was very recently described by Crous *et al.* (2007) on leaves of *Eucalyptus* collected in Italy. Very little is known regarding this fungus but Perez *et al.* (2009b) found it on *E. dunnii*, *Eucalyptus globulus* and *Eucalyptus grandis* in the northern region of Uruguay. Although *Pseudocercospora luzardii* grouped closely with *Ps. norchiensis*, the similarity in morphological features of isolates UY1436 and UY1484 and those described for *Ps. norchiensis* as well as the DNA sequence data (100 % of similarity with the ex-type sequence of *Ps. norchiensis*) supports the identification of these isolates as *Ps. norchiensis*. In addition, the reference sequence of *Ps. luzardii* (AF362057) in GenBank differed from *Ps. norchiensis* (EF394859) at three nucleotides in the ITS2 region. The former species has been reported only on *Hancornia speciosa* (Apocynaceae) in Brazil (Furnaletto & Dianese 1999) and it probably represents a distinct species.

Mycosphaerella yunnanensis found on native Myrtaceae in this study has been described by Burgess *et al.* (2007) from *Eucalyptus urophylla* in China. We found this species associated with leaf spots on the native *B. salicifolius* and *Myrrhinium atropurpureum* var. *octandrum* in Uruguay. To the best of our knowledge, this is the first report of *M. yunnanensis* outside China. Although it has not been found on *Eucalyptus* in Uruguay, it seems likely that its origin is on that host.

Passalora loranthei appears to be a species in the Mycosphaerellaceae with a wide host range. The fungus has been previously recorded in two unrelated hosts, namely *Citrus* sp. and *Musa* (Arzanlou *et al.* 2008). We found *Pa. loranthei* associated with leaf disease on *A. sellowiana*. This finding adds a member of the Myrtaceae to the list of hosts that can be infected by this fungus.

This study is the first to broadly consider the MLD on native Myrtaceae growing in association with non-native *Eucalyptus* plantations. Four species previously only known from *Eucalyptus* were found on native Myrtaceae. This suggests strongly that these fungi are moving from non-native *Eucalyptus* to native trees. Our findings support the views of Crous & Groenewald (2005) and Hunter *et al.* (2011) that some leaf-infecting fungi previously thought to be specific to *Eucalyptus*, have wider host ranges than was thought in the past. Almost nothing is known regarding the etiology and impact of these species, other than the fact that they are associated with leaf spot diseases on native trees. There is currently no evidence to suggest that they are causing serious disease problems on the native trees on which they were found, but their potential to result in serious disease situations such as those observed on *Eucalyptus* in Uruguay and around the world must be considered. Continued monitoring of these disease situations is essential.

While there are growing numbers of examples of pathogens of native Myrtaceae moving to *Eucalyptus* where these trees are grown as exotics (Rodas *et al.* 2005; Glen *et al.* 2007; Coutinho *et al.* 2011), there are far fewer examples of movement of apparently introduced *Eucalyptus* pathogens to native plants (Crous & Groenewald 2005). Results of this study provide important new information that this movement is far more common than has been previously thought. Although the consequences have yet to be realized, the results illustrate the danger of moving crop plants between

countries together with pathogenic fungi that are poorly understood.

Acknowledgements

We are grateful to Forestal Oriental, Rivermol, Stora Enso and Weyerhaeuser for providing funds to support this study. We appreciate the assistance of Andrés Berrutti, Ana Terzaghi, Sofía Simeto and Gustavo Balmelli on this work.

REFERENCES

- Andjic V, Whyte G, Hardy G, Burgess, 2010. New *Teratosphaeria* species occurring on eucalypts in Australia. *Fungal Diversity* 43: 27–38.
- Arzanlou M, Groenewald JZ, Fullerton RA, Abeln ECA, Carlier J, Zapater MF, Buddenhagen IW, Viljoen A, Crous PW, 2008. Multiple gene genealogies and phenotypic characters differentiate several novel species of *Mycosphaerella* and related anamorphs on banana. *Persoonia* 20: 19–37.
- Balmelli G, Marroni V, Altier N, García R, 2004. Potencial del mejoramiento genético para el manejo de enfermedades en *Eucalyptus globulus*. In: *Serie Técnica* 143. INIA, Montevideo, Uruguay.
- Brussa CA, Grela IA, 2007. *Flora arbórea del Uruguay con énfasis en las especies de Rivera y Tacuarembó*. COFUSA, Mosca. Montevideo, Uruguay.
- Burgess TI, Barber PA, Sufaati S, Xu D, Hardy GESTJ, Dell B, 2007. *Mycosphaerella* spp. on *Eucalyptus* in Asia; new species, new hosts and new records. *Fungal Diversity* 24: 135–157.
- Burgess T, Wingfield MJ, 2002. Quarantine is important in restricting the spread of exotic seed-borne tree pathogens in the southern hemisphere. *International Forestry Review* 4: 56–65.
- Carnegie AJ, Ades PK, Keane PJ, Smith IW, 1998. *Mycosphaerella* diseases of juvenile foliage in a eucalypt species and provenance trial in Victoria. Australia. *Australian Forestry* 61: 190–194.
- Carnegie AJ, Burgess TI, Beilharz V, Wingfield MJ, 2007. New species of *Mycosphaerella* from Myrtaceae in plantations and native forest in Australia. *Mycologia* 99: 461–474.
- Carnegie AJ, Keane PJ, Ades PK, Smith IW, 1994. Variation in susceptibility of *Eucalyptus globulus* provenances to *Mycosphaerella* leaf disease. *Canadian Journal of Forest Research* 24: 1751–1757.
- Carnegie AJ, Lidbetter JR, Walker J, Horwood MA, Tesoriero L, Glen M, Priest MJ, 2010. *Uredo rangelii*, a taxon in the guava rust complex, newly recorded on Myrtaceae in Australia. *Australasian Plant Pathology* 39: 463–466.
- Carbone I, Kohn LM, 1999. A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia* 91: 553–556.
- Cortinas MN, Crous PW, Wingfield BD, Wingfield MJ, 2006. Multi-gene phylogenies and phenotypic characters distinguish two species within the *Colletogloeopsis zuluensis* complex associated with *Eucalyptus* stem cankers. *Studies in Mycology* 55: 113–146.
- Coutinho TA, Brady CL, van der Vaart MM, Venter SN, Telechea N, Rolfo M, Pérez C, Wingfield MJ, 2011. A new shoot and stem disease of *Eucalyptus* species caused by *Erwinia psidii*. *Australasian Plant Pathology* 40: 55–60.
- Coutinho TA, Wingfield MJ, Alfenas AC, Crous PW, 1998. *Eucalyptus* rust: a disease with the potential for serious international implications. *Plant Disease* 82: 819–825.
- Crous PW, 1998. *Mycosphaerella* spp. and Their Anamorphs Associated with Leaf Spot Diseases of *Eucalyptus*. APS Press, St. Paul, Minnesota, USA.

- Crous PW, 1999. Species of *Mycosphaerella* and related anamorphs occurring on Myrtaceae (excluding *Eucalyptus*). *Mycological Research* **103**: 607–621.
- Crous PW, Braun U, Hunter GC, Wingfield MJ, Verkley GJM, Shin HD, Nakashima C, Groenewald JZ, 2013. Phylogenetic lineages in *Pseudocercospora*. *Studies in Mycology* **75**: 37–114.
- Crous PW, Groenewald JZ, 2005. Hosts, species and genotypes: opinions versus data. *Australasian Plant Pathology* **34**: 463–470.
- Crous PW, Groenewald JZ, Mansilla JP, Hunter GC, Wingfield MJ, 2004. Phylogenetic reassessment of *Mycosphaerella* spp. and their anamorphs occurring on *Eucalyptus*. *Studies in Mycology* **50**: 195–214.
- Crous PW, Summerell BA, Carnegie AJ, Mohammed C, Himaman W, Groenewald JZ, 2007. Foliicolous *Mycosphaerella* spp. and their anamorphs on *Corymbia* and *Eucalyptus*. *Fungal Diversity* **26**: 143–185.
- Crous PW, Summerell BA, Carnegie AJ, Wingfield MJ, Groenewald JZ, 2009. Novel species of *Mycosphaerellaceae* and *Teratosphaeriaceae*. *Persoonia* **23**: 119–146.
- Crous PW, Wingfield MJ, Mansilla JP, Alfenas AC, Groenewald JZ, 2006. Phylogenetic reassessment of *Mycosphaerella* spp. and their anamorphs occurring on *Eucalyptus* II. *Studies in Mycology* **55**: 99–131.
- Farris JS, Källersjö M, Kluge AG, Bult C, 1995. Testing significance of incongruence. *Cladistics* **10**: 315–319.
- Furnaletto C, Dianese JC, 1999. Some *Pseudocercospora* species and new *Prathigada* species from the Brazilian cerrado. *Mycological Research* **103**: 1203–1209.
- Glen M, Alfenas AC, Zauza EAV, Wingfield MJ, Mohammed C, 2007. *Puccinia psidii*: a threat to the Australian environment and economy – a review. *Australasian Plant Pathology* **36**: 1–16.
- Grgurinovic CA, Walsh D, Macbeth F, 2006. *Eucalyptus* rust caused by *Puccinia psidii* and the threat it poses to Australia. *EPPO Bulletin* **36**: 486–489.
- Gryzenhout M, Rodas CA, Mena Portales J, Clegg P, Wingfield BD, Wingfield MJ, 2006. Novel hosts of the *Eucalyptus* canker pathogen *Chrysosporthe cubensis* and a new *Chrysosporthe* species from Colombia. *Mycological Research* **110**: 833–845.
- Heath RN, Gryzenhout M, Roux J, Wingfield MJ, 2006. Discovery of the canker pathogen *Chrysosporthe austroafricana* on native *Syzygium* spp. in South Africa. *Plant Disease* **90**: 433–438.
- Hillis DM, Bull JJ, 1993. An empirical test of bootstrapping as a method for assessing confidence in phylogenetic analysis. *Systematic Biology* **42**: 142–152.
- Hodges CS, Alfenas AC, Cordell CE, 1986. The conspecificity of *Cryphonectria cubensis* and *Endothia eugeniae*. *Mycologia* **78**: 334–350.
- Huelsbeck JP, Bull JJ, Cunningham CW, 1996. Combining data in phylogenetic analysis. *TREE* **11**: 152–158.
- Hunter GC, Crous PW, Carnegie AJ, Burgess TI, Wingfield MJ, 2011. *Mycosphaerella* and *Teratosphaeria* diseases of *Eucalyptus*; easily confused and with serious consequences. *Fungal Diversity* **50**: 145–166.
- Hunter GC, Roux J, Wingfield BD, Crous PW, Wingfield MJ, 2004. *Mycosphaerella* species causing leaf disease in South African *Eucalyptus* plantations. *Mycological Research* **108**: 672–681.
- Hunter GC, Wingfield BD, Crous PW, Wingfield MJ, 2006. Multi-gene phylogeny for *Mycosphaerella* species occurring on *Eucalyptus* leaves. *Studies in Mycology* **55**: 147–161.
- Katoh K, Kuma K, Toh H, Miyata T, 2005. MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Research* **33**: 511–518.
- Lundquist JE, Purnell RC, 1987. Effects of *Mycosphaerella* leaf spot on growth of *Eucalyptus nitens*. *Plant Disease* **71**: 1025–1029.
- Maxwell A, Dell B, Neumeister-Kemp H, Hardy GESTJ, 2003. *Mycosphaerella* species associated with *Eucalyptus* in south-western Australia: new species, new records and a key. *Mycological Research* **107**: 351–359.
- Myburg H, Gryzenhout M, Wingfield BD, Wingfield MJ, 2003. Conspecificity of *Endothia eugeniae* and *Cryphonectria cubensis*: a re-evaluation based on morphology and DNA sequence data. *Mycoscience* **104**: 187–196.
- Park RF, Keane PJ, Wingfield MJ, Crous PW, 2000. Fungal diseases of eucalypt foliage. In: Keane PJ, Kile GA, Podger FD, Brown BN (eds), *Diseases of Pathogens of Eucalypts*. CSIRO Publishing, Melbourne, pp. 153–240.
- Perez CA, de Beer ZW, Altier NA, Wingfield MJ, Blanchette RA, 2008. Discovery of the eucalypt pathogen *Quambalaria eucalypti* infecting a non-*Eucalyptus* host in Uruguay. *Australasian Plant Pathology* **37**: 600–604.
- Perez G, Hunter GC, Slippers B, Perez C, Wingfield BD, Wingfield MJ, 2009a. *Teratosphaeria* (*Mycosphaerella*) *nubilosa*, the causal agent of *Mycosphaerella* Leaf Disease (MLD), recently introduced into Uruguay. *European Journal of Plant Pathology* **125**: 109–118.
- Perez C, Wingfield MJ, Altier NA, Blanchette RA, 2009b. *Mycosphaerellaceae* and *Teratosphaeriaceae* associated with *Eucalyptus* leaf diseases and stem cankers in Uruguay. *Forest Pathology* **39**: 349–360.
- Perez C, Wingfield MJ, Slippers B, Altier NA, Blanchette RA, 2009c. *Neofusicoccum eucalyptorum*, a *Eucalyptus* pathogen, on native Myrtaceae in Uruguay. *Plant Pathology* **58**: 964–970.
- Posada D, Crandall KA, 1998. MODELTEST: testing the model of DNA substitution. *Bioinformatics* **14**: 817–818.
- Rodas CA, Gryzenhout M, Myburg H, Wingfield BD, Wingfield MJ, 2005. Discovery of the *Eucalyptus* canker pathogen *Cryphonectria cubensis* on native *Miconia* (*Melastomataceae*) in Colombia. *Plant Pathology* **54**: 460–470.
- Sánchez Márquez S, Bills GF, Zabalgoceazcoa I, 2011. Fungal species diversity in juvenile and adult leaves of *Eucalyptus globulus* from plantations affected by *Mycosphaerella* leaf disease. *Annals of Applied Biology* **158**: 177–187.
- Sivanesan A, Shivas RG, 2002. Studies on *Mycosphaerella* species in Queensland, Australia. *Mycological Research* **106**: 355–364.
- Slippers B, Stenlid J, Wingfield MJ, 2005. Emerging pathogens: fungal host jumps following anthropogenic introduction. *Trends in Ecology and Evolution* **20**: 420–421.
- Summerell BA, Groenewald JZ, Carnegie A, Summerbell RC, Crous PW, 2006. *Eucalyptus* microfungi known from culture 2. *Alysidiella*, *Fusculina* and *Phlogicylindrium* genera nova, with notes on some other poorly known taxa. *Fungal Diversity* **23**: 323–350.
- Swofford DL, 2002. PAUP*: *phylogenetic analysis using parsimony (*and other methods)*. Version 4.0b10a. Sinauer Associates, Sunderland, MA.
- White TJ, Bruns S, Lee S, Taylor JW, 1990. Amplification and direct sequencing of fungal genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (eds), *PCR Protocols: a guide to methods and applications*. Academic Press, San Diego, pp. 315–322.
- Wingfield MJ, 2003. Increasing threat of disease to exotic plantation forests in the Southern Hemisphere: lessons from *Cryphonectria* canker. *Australasian Plant Pathology* **32**: 133–139.
- Wingfield MJ, Slippers B, Hurley BP, Coutinho TA, Wingfield BD, Roux J, 2008. *Eucalypt* pests and diseases: growing threats to plantation productivity. *Southern Forests* **70**: 139–144.